

Two Anderson Impurities in the Kondo Limit Doublet States

J. Simonin

*Centro Atómico Bariloche and Instituto Balseiro,
8400 S.C. de Bariloche, Río Negro, Argentina*

(Dated: mar)

We analyze two pairs of doublet states for the two Anderson impurity problem. We found that for short interimpurity distances they have a lower energy than the ferro triplet and the antiferro singlet. For medium to long distances between the impurities, the doublets also have a lower energy than the two decoupled Kondo-singlet state for most of the range of interest. The mechanism behind their energy gain is an coherence enhanced Kondo-like interaction.

PACS numbers: 73.23.-b, 72.15.Qm, 73.63.Kv, 72.10.Fk

The two Anderson impurity problem, i.e. how two magnetic impurities interact when they are embedded in a metallic host, has been the subject of many theoretical studies, ranging from perturbation theory^{1,2,3} and narrow band approximation⁴ to renormalization group analysis⁵. Besides its natural presence in alloys, today this system can be tailor made by means of quantum dots^{6,7} and promises to be a relevant circuit component in quantum electronics. Here we present a variational wave function analysis that, as in the single impurity case⁸, gives a clear description of the system. When needed, we specialize in the one dimensional (1D) case because of its technological applications.

The Anderson Hamiltonian for magnetic impurities diluted in a metallic host is:

$$H = \sum_{k\sigma} e_k c_{k\sigma}^\dagger c_{k\sigma} + \frac{V}{\sqrt{N_c}} \sum_{jk\sigma} (e^{i\mathbf{k}\cdot\mathbf{r}_j} d_{j\sigma}^\dagger c_{k\sigma} + h.c.) - E_d \sum_{j\sigma} d_{j\sigma}^\dagger d_{j\sigma} + U \sum_j d_{j\uparrow}^\dagger d_{j\uparrow} d_{j\downarrow}^\dagger d_{j\downarrow} , \quad (1)$$

where the fermion operators $c_{k\sigma}$ act on the conduction band states and $d_{j\sigma}$ on the orbital of the magnetic impurity situated at $\mathbf{r}_j$. Single state energies $e_k, -E_d$ are referred to the Fermi energy ($E_F = k_F^2/2m$), i.e. there is an implicit $-\mu N$ term in the Hamiltonian that regulates the population of the system ($\mu = E_F$ is the chemical potential and N the total number operator), V is the $d-c$ hybridization and N_c is the number of cells in the metal. In the Kondo limit the impurity levels are well below the Fermi energy ($E_d \gg 0$), and they can not be double occupied due to the Coulomb repulsion in them ($U \gg 2E_d$). In order to simplify calculations we renormalize the vacuum (denoted by $|F\rangle$) to be the conduction band filled up to the Fermi energy and we make an electron-hole transformation for band states below the Fermi level: $b_{k\sigma}^\dagger \equiv c_{-k,-\sigma}$ for $|k| \leq k_F$. In this way the energy of a hole excitation (i.e., to remove an electron from below the Fermi level) is explicitly positive.

We consider here the two impurities case, one impurity at $-\mathbf{r}/2$ (the Left impurity) and the other at $\mathbf{r}/2$ (the Right impurity). Although calculations were made with the second quantization operators, we use in the text a

“ket” notation for the impurity states: the first symbol indicates the population status of the Left impurity and the second the status of the Right impurity, e.g. $|\downarrow, \uparrow\rangle \equiv d_{L\downarrow}^\dagger d_{R\uparrow}^\dagger |F\rangle$, $|0, \uparrow\rangle \equiv d_{R\uparrow}^\dagger |F\rangle$, $|0, 0\rangle \equiv |F\rangle$, etc..

In this situation ($E_d \gg 0$, $U \gg 2E_d$) the lower energy configurations of the system are the ferro triplet ($|\uparrow, \uparrow\rangle$, $(|\uparrow, \downarrow\rangle + |\downarrow, \uparrow\rangle)$, $|\downarrow, \downarrow\rangle$) and the antiferro singlet ($(|\uparrow, \downarrow\rangle - |\downarrow, \uparrow\rangle)$), with energies $-2E_d$. These states are not eigenstates of the system; the hybridization mixes these states with configurations having excitations in the band. Perturbation theory allows for the calculation of the corresponding corrections. These energy corrections are the same for both the ferro (FF) and antiferro (AF) states (and twice the single impurity corrections) up to the well known RKKY-like four order term, which has opposite contributions on them¹²:

$$E_{FF} = -2E_d - \Sigma_R , \quad E_{AF} = -2E_d + \Sigma_R , \quad (2)$$

$$\Sigma_R = 2 \mathbf{v}^4 \sum_{k,q} \frac{\cos((\mathbf{q} + \mathbf{k})\cdot\mathbf{r})}{(E_d + e_k)^2 (e_k + e_q)} , \quad (3)$$

where the q (k) sum is over hole (electron) excitations ($|q| \leq k_F$, $|k| \geq k_F$) and $\mathbf{v} = V/\sqrt{N_c}$.

A related state is the Kondo singlet, the ground state when just one impurity is considered. If only the Left impurity is present, this singlet is described by the variational wave function⁸

$$|S_L\rangle = |F\rangle - \sum_q \alpha_q^K e^{i\mathbf{q}\cdot\mathbf{r}/2} (b_{q\uparrow}^\dagger |\downarrow_L\rangle + b_{q\downarrow}^\dagger |\uparrow_L\rangle) , \quad (4)$$

where the variational parameter results to be $\alpha_q^K = \mathbf{v}/(E_S + E_d - e_q)$ and a self-consistent equation is obtained for the singlet energy E_S ,

$$E_S = 2 \mathbf{v}^2 \sum_q \frac{1}{E_S + E_d - e_q} , \quad (5)$$

after the pole in the sum, $E_S = -E_d - \delta_K$ is proposed ($E_d \gg \delta_K$), and Eq.(5) becomes an equation for the Kondo energy δ_K ,

$$E_d = 2 \mathbf{v}^2 \sum_q \frac{1}{\delta_K + e_q} , \quad (6)$$

from which

$$\delta_K = D e^{-1/(2Jn)} \quad (7)$$

is obtained. Jn ($\equiv (V^2/E_d) n_o$, n_o being the density of states at the Fermi level) is the relevant parameter for these theories, and D , the half band-width, becomes the energy scale (δ_K , Σ_R are proportional to it). $Jn = 0.07238$ gives δ_K one thousandth of D , a reasonable value.

We study two doublet states, the even doublet,

$$\begin{aligned} |D_E\rangle = & (|\uparrow, 0\rangle + |0, \uparrow\rangle) + \sum_q \{ \\ & -i \alpha_q^F \sin \frac{q.r}{2} [2b_{q\downarrow}^\dagger |\uparrow, \uparrow\rangle + b_{q\uparrow}^\dagger (|\uparrow, \downarrow\rangle + |\downarrow, \uparrow\rangle)] \\ & + \alpha_q^A \cos \frac{q.r}{2} b_{q\uparrow}^\dagger (|\uparrow, \downarrow\rangle - |\downarrow, \uparrow\rangle) \} , \quad (8) \end{aligned}$$

and the odd doublet,

$$\begin{aligned} |D_O\rangle = & (|\uparrow, 0\rangle - |0, \uparrow\rangle) + \sum_q \{ \\ & + \alpha_q^F \cos \frac{q.r}{2} [2b_{q\downarrow}^\dagger |\uparrow, \uparrow\rangle + b_{q\uparrow}^\dagger (|\uparrow, \downarrow\rangle + |\downarrow, \uparrow\rangle)] \\ & -i \alpha_q^A \sin \frac{q.r}{2} b_{q\uparrow}^\dagger (|\uparrow, \downarrow\rangle - |\downarrow, \uparrow\rangle) \} . \quad (9) \end{aligned}$$

After a standard minimization procedure, the variational parameters are obtained

$$\alpha_q^{F(A)} = \frac{\mathbf{v}}{E_X + 2E_d \pm \Sigma_R - e_q} , \quad (10)$$

where E_X stands for the energy of the doublet (E_E or E_O) and the plus sign in front of Σ_R holds for the ferro α factor (α^F) and the minus for the antiferro. For the energies, self-consistent equations result

$$\begin{aligned} E_E = & -E_d + \mathbf{v}^2 \sum_q \frac{3 \sin^2 \frac{q.r}{2}}{E_E + 2E_d + \Sigma_R - e_q} \\ & + \mathbf{v}^2 \sum_q \frac{\cos^2 \frac{q.r}{2}}{E_E + 2E_d - \Sigma_R - e_q} . \quad (11) \end{aligned}$$

For E_O a similar equation holds, with the sin and cos interchanged. This behavior, as well as the sign of Σ_R , is due to the spatial symmetry of the ferro and antiferro states. As is the case of the Kondo singlet, the dominant pole in the sums of Eq.(11) determines the value of E_E .

First, let us analyze the situation when Σ_R vanishes. This situation corresponds to the zeros of the RKKY interaction and the $k_{Fr} \gg \pi$ limit.

For $\Sigma_R = 0$, after expanding the $\sin^2$, $\cos^2$ functions, Eq.(11) reads

$$E_E = -E_d + \mathbf{v}^2 \sum_q \frac{2 - \cos q.r}{E_E + 2E_d - e_q} . \quad (12)$$

Proposing $E_E = -2E_d - \delta_E$ we obtain

$$E_d = \mathbf{v}^2 \sum_q \frac{2 - \cos q.r}{\delta_E + e_q} . \quad (13)$$

Therefore for very large r , such that the contribution from the cos term vanishes, the energy gain of the doublet (δ_E) tends to that of one Kondo singlet. This is not surprising, given that the doublets, when $\Sigma_R = 0$, can be written as

$$|D_X\rangle = |\uparrow_L\rangle \otimes |S_R\rangle \pm |S_L\rangle \otimes |\uparrow_R\rangle, \quad (14)$$

i.e., the combination of a Kondo singlet in one impurity and the other single occupied plus (minus) their mirror image.

This energy, δ_E , must be compared with twice the Kondo energy, because, for very large r , the latter is the energy gain for a state that has simultaneously both impurities forming a Kondo singlet ($|S_L\rangle \otimes |S_R\rangle$).

Now, we examine the effect of the coherence term in Eq.(13); here we present the 1D analysis. The sums in Eq.(13) can be done exactly. The first one is the Kondo integral and we call the second one the Doublet Coherence integral,

$$\frac{1}{N_c} \sum_q \frac{1}{\delta + e_q} = n_o I_K(\delta) = n_o \ln(1 + \frac{D}{\delta}) , \quad (15)$$

$$\frac{1}{N_c} \sum_q \frac{\cos q.r}{\delta + e_q} = n_o I_X(\delta, x) , \quad (16)$$

where $x = k_{Fr}$. The Coherence integral has a logarithmic dependence on δ that goes like I_K , so it is useful to define $C_D(\delta, x) = I_X(\delta, x)/I_K(\delta)$, which is a decaying oscillatory function that depends lightly on δ . We will see that C_D plays the role of a coherence driven extra connectivity between the components of the doublet wave function. In Fig.1 we plot C_D as a function of k_{Fr} for various values of δ , and the r dependence of the 1D-RKKY is also plotted. It can be seen that the RKKY reaches its asymptotic form ($-\cos(2x)/\pi x$) very quickly, at $k_{Fr} \geq \pi$, whereas I_X approaches a similar form ($\sin(x) D/x\delta$) but at $k_{Fr} \delta/D \geq \pi$, a distance about one thousand times greater, the Kondo length.

As a function of I_K and C_D Eq.(13) becomes

$$1/Jn = [2 - C_D(\delta_E, k_{Fr})] I_K(\delta_E) , \quad (17)$$

from which δ_E can be obtained numerically. It is interesting to write it the Kondo way (Eq.(7)),

$$\delta_E = D \exp \left[\frac{-1}{[2 - C_D(\delta_E, k_{Fr})] Jn} \right] , \quad (18)$$

which can be taken as an approximate solution by evaluating C_D at δ_K , or used iteratively to find the exact result with fast convergency. Note that C_D couples to the ‘‘connectivity’’ factor (the 2 in the exponent of Eq.(7)), and small changes in this factor produce huge changes in δ_E . For δ_O the sign in front of C_D is positive. These energies are plotted in Fig.2 as a function of k_{Fr} for $\delta_K = 0.001 D$. They alternatively surpass $2 \delta_K$ up to $k_{Fr} \simeq 50\pi$.

This effect, $\delta_E, \delta_O \geq 2\delta_K$, is present for values of Jn lower than $Jn_c = 0.240$ (which gives $\delta_K = D/8$); the




FIG. 1: Range functions for the RKKY, $J_R(2k_F r)$, and the Doublet Coherence factor, $C_D(\delta, k_F r)$. The period of the RKKY is half the one of C_D and its amplitude decays very rapidly. These characteristics are due to the fact that the RKKY depends on two excitations, whereas the Doublet Coherence involves just one.




FIG. 2: Correlation energy gains of the odd, δ_O , and even, δ_E , doublets as a function of the distance between the impurities, for $\Sigma_R = 0$ and $Jn = 0.07238$. The inset is δ_E up to very long distances, and it periodically beats $2\delta_K$ up to $k_F r \simeq 50\pi$.

lower Jn , the greater this effect. The periodicity with which $|D_O\rangle$, $|D_E\rangle$ alternate is approximately twice the RKKY period (they are ruled by C_D), and they stand in this situation up to very long distances. These properties are due to the fact that the Doublet Coherence integral involves just one hole excitation. At the intersection points one has $\delta_E, \delta_O = \delta_K$ and $C_D = 0$.

The situation for $\Sigma_R \neq 0$ can be analyzed with the same concepts already introduced. We analyze here the even doublet energy E_E when $\Sigma_R > 0$. The analysis for E_O and/or $\Sigma_R < 0$ is very similar.

For $\Sigma_R > 0$ the lower energy configurations are the ones of the ferro triplet, which makes the pole of α_q^F the relevant one in the equation for E_E , Eq.(11). Thus,

proposing $E_E = -2E_d - \Sigma_R - \delta_E$, Eq.(11) is transformed to

$$E_d + \Sigma_R + \delta_E = \frac{3}{2} v^2 \sum_q \frac{1 - \cos qr}{\delta_E + e_q} + \frac{1}{2} v^2 \sum_q \frac{1 + \cos qr}{2\Sigma_R + \delta_E + e_q}, \quad (19)$$

and then, using the definitions for I_K and I_X , we have,

$$\frac{E_d + \Sigma_R + \delta_E}{n_o V^2} = \frac{3}{2} [I_K(\delta_E) - I_X(\delta_E)] + \frac{1}{2} [I_K(2\Sigma_R + \delta_E) + I_X(2\Sigma_R + \delta_E)], \quad (20)$$

a self-consistent equation for δ_E (in Eq.(20) I_X is also a function of $k_F r$, but only the “energy” dependence is quoted). If $E_d \gg \Sigma_R, \delta_E$ the left side of Eq.(20) is just $1/Jn$; if not, a new parameter must be introduced: the E_d/D ratio. In the present work we keep it simple (just Jn). For δ_O (and $\Sigma_R > 0$) the sign of the I_X terms are inverted. Therefore when δ_O decreases δ_E increases and viceversa, following the oscillations of C_D as a function of $k_F r$, just as in the $\Sigma_R = 0$ case. For $\Sigma_R < 0$ the dominant pole is that of the antiferro component of the doublets and δ_E, δ_O swap equations, with $|\Sigma_R|$ instead of Σ_R and one important difference: the bare connectivity of the antiferro component is $1/2$, instead of the $3/2$ of the ferro one, and hence these energy gains are lower than in the $\Sigma_R > 0$ case.




FIG. 3: Total energy gain of the even doublet, $\Delta_E = |\Sigma_R| + \delta_E$, as a function of r . Note that in the relevant ranges for this state the main contribution to Δ_E comes from δ_E .

In Fig.3 we show δ_E , Σ_R , and the total energy gain of the even doublet, $\Delta_E = |\Sigma_R| + \delta_E$. Note the kinks in δ_E at the zeros of Σ_R , those kinks reflect the change of dominant component in $|D_E\rangle$ when Σ_R changes sign. Nevertheless, the total energy gain is smooth at those points. In the regions in which δ_E is equal to zero the doublet “collapses”: i.e. when $\delta_E = 0$ the dominant $\alpha_q^{F(A)}$ diverges for $|q| = k_F$, (Eqs.(8,10), $e_{k_F} = 0$) and

thus this is the only surviving configuration of the ones that compose the doublet, and a FF (AF) state with just one hole at the Fermi level is simply a FF (AF) state. The kinks of Δ_E in these regions are because $\Delta_E = |\Sigma_R|$. In the relevant regions ($\delta_E > \delta_O$) usually $\delta_E > |\Sigma_R|$, thus the dominant pole configurations (FF (AF) based) have just a little more weight in the doublet than the “dominated” one (AF (FF) based), and the $\Sigma_R = 0$ regime is soon reached.

Finally, in Fig.4, we plot both Δ_E and Δ_O as function of r , Σ_R is also shown ($\delta_K = 0.001D$). For this value of Jn , in the measure that r increases the system alternates between the odd and even doublets, just as in the $\Sigma_R = 0$ case, except for a small region above $k_F r = \pi/2$. At this point Σ_R is near an AF maximum and C_D is at its first zero. Hence there is not connectivity enhancement, neither for the odd, nor for the even doublet, and they collapse into the AF state.




FIG. 4: Δ_E and Δ_O as a function of r , for $\delta_K = 0.001D$.

In conclusion, we give a clear description of the behav-

ior of these doublet states of the Two Anderson Impurity System and their interplay with the RKKY interaction. A remarkable point of the model is the long distances at which the impurities remain in a correlated state. A full understanding of this property, which our model provides in detail, allows these systems to be used in the design of quantum devices.

I am grateful to Blas Alascio for insightful inquiries. I am a fellow of the CONICET (Argentina), which partially financed this research under grant PIP 02753/00.

Appendix: In 1D, Σ_R , Eq.(2), half the RKKY interaction, is

$$\begin{aligned} \Sigma_R &\simeq 2 \frac{v^4}{E_d^2} \sum_{k,q} \frac{\cos(q.r) \cos(k.r)}{(e_k + e_q)} \\ &= \left(\frac{\pi}{2} Jn\right)^2 D J_R(2k_F r), \end{aligned} \quad (\text{A.21})$$

where the r dependence is given by⁹,

$$J_R(2k_F r) = \frac{2}{\pi} \left[\frac{\pi}{2} - \text{Si}(2k_F r) \right]. \quad (\text{A.22})$$

The 1D Doublet Coherence integral is given by

$$\begin{aligned} I_X(\delta, x) = & \cos(x + xd) [\text{Ci}(x + xd) - \text{Ci}(x)] + \\ & \sin(x + xd) [\text{Si}(x + xd) - \text{Si}(x)], \end{aligned} \quad (\text{A.23})$$

where $d = \delta/D$. Ci (Si) is the CosIntegral (SinIntegral) function, as defined in Mathematica[®].

The upper critical value of Jn , given that $|C_D| \leq 1$, is defined by the equation

$$e^{-1/(3Jn_c)} = 2 e^{-1/(2Jn_c)}, \quad (\text{A.24})$$

which gives $Jn_c = 1/6 \ln 2$.

-
- ¹ B. Coqblin and J. R. Schieffer, Phys. Rev. **185**, 847 (1969).
² C. R. Proetto and A. López, Phys. Rev. B **25**, 7037 (1982).
³ S. Galkin, C. A. Balseiro, and M. Avignon, Euro. Phys. J. B **38**, 519 (2004).
⁴ R. Allub, Phys. Rev. B **67**, 144416 (2003).
⁵ B. A. Jones and C. M. Varma, Phys. Rev. Lett. **58**, 843 (1987).
⁶ N. J. Craig, J. M. Taylor, E. A. Lester, C. M. Marcus, M. P.

- Hanson, and A. C. Gossard, Science **304**, 565 (2004).
⁷ P. Simon (2004), preprint: Probing the Kondo screening cloud in a double quantum dot set up.
⁸ C. M. Varma and Y. Yafet, Phys. Rev. B **13**, 2950 (1976).
⁹ V. I. Litvinov and V. K. Dugaev, Phys. Rev. B **58**, 3584 (1998).